



A tropical tale: how *Naja nigricollis* venom beats *Trypanosoma brucei*

Martos Esteban, Andrea; Laustsen, Andreas Hougaard; Carrington, Mark

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):


Martos Esteban, A., Laustsen, A. H., & Carrington, M. (2016). *A tropical tale: how Naja nigricollis venom beats Trypanosoma brucei*. Poster session presented at Symposium for Biological and Life Science Students 2016, Cambridge, United Kingdom.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



A tropical tale: how *Naja nigricollis* venom beats *Trypanosoma brucei*


Andrea Martos Esteban¹, Andreas Hougaard Laustsen¹, Mark Carrington²

¹Department of Biotechnology and Biomedicine, Technical University of Denmark

²Department of Biochemistry, University of Cambridge

Sleeping sickness: a neglected tropical disease


Trypanosoma brucei is a parasitic protozoan species capable to infecting insect vectors whose bite further produces African sleeping sickness in human beings [1]. During the parasite's extracellular life in the mammalian host, its outer coat, mainly composed of Variable Surface Glycoproteins (VSGs) [2], undergoes enormous variation in its composition to avoid the host's lytic immune response [3].



N. nigricollis venom is able to kill *T. brucei* by targeting GPI anchoring of VSG


Phospholipases A₂ are one of the major enzymatically active components in *N. nigricollis* venom, which could be targeting the GPI anchor of VSG. Lyophilized whole venom was diluted in HMI-9 cell culture media, in which we cultured *T. brucei* parasites expressing eGFP attached to GPI (VSG121) with 10 µm/ml of *N. nigricollis* venom during 24 hours. Then, we study the surveillance and the GFP-GPI release by flow cytometry.

Parasite cells counts by flow cytometry



PLA₂ and cytotoxins are potentially the toxins responsible of the parasite death

After gel filtration of 20 mg of *N. nigricollis* whole venom, we tested each venom fraction in culture and ran a SDS-PAGE, which was developed by silver-staining.




References

- [1] Kennedy PG E., 2013. The Lancet. DOI:10.1016/S1474-4422(12)70296-X.
- [2] Sunter J., et al. 2013. PLOS Pathogen. DOI:10.1371/journal.ppat.1003566.
- [3] Fenn K. and Matthews K R., 2007. Current Opinion in Microbiology. DOI:10.1016/j.mib.2007.09.014.
- [4] Petras D., et al. 2011. Journal of Proteome Research. DOI:10.1021/pr101040f


A tropical tale of trypanosomes and snakes

The elapid *N. nigricollis* is a large venomous snake from sub-Saharan Africa [4]. Since the origin of pharmacology, venoms have been used as medicines, since venom toxins target a myriad of different physiological processes with high specificity and selectivity.



N. nigricollis venom provokes the flagellar pocket enlargement by accumulating GPI

We observed VSG121 parasites using the fluorescence microscope. Contrarily to the parasites' phenotype during venom absence, parasites cultured with the *N. nigricollis* venom show a greener and enlarged flagellar pocket over the time.



Impact

I. First report of using snake venom to effectively kill effectively *T. brucei*.

II. Novel molecular target antigenic variation-independent, allowing to bypass the main challenge of developing a successful treatment for the neglected tropical disease, African sleeping sickness.

III. Unveiling of the mechanism of parasite lethality may help pave the way for novel molecular tools for drug discovery against trypanosome related diseases.



Contact information

DTU Biotechnology and Biomedicine
Technical University of Denmark
Søltofts Plads 223 – 2800 – Kgs. Lyngby, Denmark
andrea.martos.esteban@gmail.com / (+34) 676 181 350

Acknowledgements

We thank the Novo Nordisk Foundation (NNF16OC0019248), Symphogen A/S and Otto Mønsted Fond for financial support.